



University of Kentucky  
**UKnowledge**

---

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII  
International Rangeland Congress

---

## The Research on the Sodium Bicarbonate Solution Cures the Ryegrass Hay

Zhilan Mu  
*Southwest University, China*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/15-2/30>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## The research on the sodium bicarbonate solution cures the ryegrass hay

Mu Zhi-Lan

Department of Animal Science ,Southwest University ,Rongchang Campus , Chongqing ,402460 . zhilanmu@yahoo .com .cn

**Key word** sodium bicarbonate solution ,drying rate ,ryegrass hay , leaf-stem ratio ,density

**Introduction** The Chongqing weather , with cloudy rainy and fog in the autumn and winter , relative humidities averaging 72% , and sometimes achieving 95% makes it very difficult to cure hay using only solar drying .Using sodium bicarbonate to destroy plant surface wax structure , promotes moisture evaporation from the plant , this increases the forage drying rate , reduces loss of leaf blades , and preserves forage nutritional value . This process also increases the leaf :stem ratio and improves smell of the hay .

**Material and method** The experimental test field is located on the Southwest University Rongchang campus . Annual precipitation averages 1099mm . Annual accumulated temperature is 6383℃ , and annual hours of sunshine total 1083h .When the weather is sunny , the temperature is 18-32℃ , the humidity 50%-70% . In the sunlight the surface temperature is 20-35℃ .

We divided samples into 5 groups stochastically , each group with 4 replications . Each perennial ryegrass sample weighed 500g . Distilled water and 0% , 0.5% , 1% , 2% , 3% sodium bicarbonate solution were sprayed on Tetragold perennial ryegrass forage samples . Samples were then exposed to the sun . After treatment were applied and samples were air dried , we determined the water content and leaf :stem ratio ,color ,smell , and ease of stem breakage .

### Result and analysis

**Table** Tetragold perennial ryegrass hay water content change and quality sense organ appraisal .

processing %	Leaf-stem ratio	Color	Ease of stem breakage	Smell	Water content (%)
0.5	1.12:1	Green shines	not easy	Light fragrant fragrance	82.26
1.0	1.14:1	Green	easy	Fragrant fragrance	80.13
2.0	1.19:1	Bottle green	easy	Rich fragrant fragrance	77.13
3.0	1.17:1	Bottle green	easy	Rich fragrant fragrance	78.67
0	1.09:1	Greyishgreen	not easy	Slightly taste	87.46

Data were analyzed using the single factor analysis of variance and multiple range test . The analysis of variance indicated the different density sprays affected the curing of perennial ryegrass hay differently ( $p < 0.01$ ) . The 2% sodium bicarbonate solution was the most effective . The data show the different solutions aided in leaf . The 2% solution significantly ( $p < 0.01$ ) increased the leaf :stem ratio 1.32% compared to the control group .

**Conclusions** Although all 4 treatment solutionsaiding in speeding up the rate of forage drying ,the result indicated that the 2% sodium bicarbonate solution was best for curing perennial ryegrass . This had the lowest water content ,highestleaf :stem ratio , and bestsmell .